

August 11, 2014

ONO PHARMACEUTICAL Co., LTD.

Corporate Communications

Mail:public_relations@ono.co.jp

**Phase 3 ASPIRE Trial of Carfilzomib (ONO-7057), a proteasome inhibitor,
in patients with relapsed multiple myeloma met
primary endpoint (Progression-free survival)**

Amgen and its subsidiary, Onyx Pharmaceuticals, Inc., (“Onyx”) announced on August 4th, 2014 (PDT) that a planned interim analysis demonstrated the Phase 3 clinical trial ASPIRE in patients with relapsed multiple myeloma met its primary endpoint of progression-free survival by administering Carfilzomib (ONO-7057, brand name in US Kyprolis[®] (carfilzomib) for Injection).

Attached is the press release distributed by Amgen and Onyx for your information.

In Japan, Ono is conducting some clinical trials in patients with multiple myeloma, in accordance with the license agreement * between Onyx and Ono signed in September 2010.

U.S. Food and Drug Administration’s (FDA) has granted accelerated approval of carfilzomib (ONO-7057), a proteasome inhibitor, indicated for the treatment of patients with multiple myeloma who have received at least two prior therapies including bortezomib and an immunomodulatory agent and have demonstrated disease progression on or within 60 days of completion of the last therapy in July 2012. Currently, no data are available for Kyprolis that demonstrate an improvement in progression-free survival or overall survival.

- * Ono entered into an exclusive license agreement with Onyx to develop and commercialize two compounds from Onyx’s proteasome inhibitor development program, carfilzomib (for injection) and oprozomib (orally administered) for all oncology indications in Japan. The compounds are both in clinical development by Onyx outside of Japan.



One Amgen Center Drive
Thousand Oaks, CA 91320-1799
Telephone 805-447-1000
www.Amgen.com

News Release

AMGEN ANNOUNCES PHASE 3 ASPIRE TRIAL OF KYPROLIS[®] IN PATIENTS WITH RELAPSED MULTIPLE MYELOMA MET PRIMARY ENDPOINT

Kyprolis Helped Patients Live 8.7 Months Longer Without Their Disease Worsening

Results to Form Basis of Regulatory Filings Beginning in 1H 2015

THOUSAND OAKS, Calif. and SOUTH SAN FRANCISCO, Calif. (Aug. 4, 2014) – Amgen (NASDAQ:AMGN) and its subsidiary, Onyx Pharmaceuticals, Inc., today announced that a planned interim analysis demonstrated that the Phase 3 clinical trial ASPIRE (CArtilizomib, Lenalidomide, and Dexamethasone versus Lenalidomide and Dexamethasone for the treatment of Patients with Relapsed Multiple Myeloma) met its primary endpoint of progression-free survival (PFS). Patients treated with Kyprolis[®] (carfilzomib) for Injection in combination with Revlimid[®] (lenalidomide) and low-dose dexamethasone (KRd) lived significantly longer without their disease worsening (median 26.3 months) compared to patients treated with Revlimid and low-dose dexamethasone (Rd) (median 17.6 months) (HR=0.690, 95 percent CI, 0.570, 0.834, $p<0.0001$). While the data for overall survival, a secondary endpoint, are not yet mature, the analysis showed a trend in favor of KRd that did not reach statistical significance.

The safety profile observed in this study is consistent with the current U.S. Kyprolis label, including the rate of cardiac events. Treatment discontinuation due to adverse events and on-study deaths were comparable between the two arms. No new safety signals were identified.

Results will be submitted for presentation at the upcoming 56th Annual Meeting of the American Society of Hematology later this year.

“We are excited about these clinical results and the positive prospects they suggest for patients with multiple myeloma,” said Robert A. Bradway, chairman and chief executive officer of Amgen, adding, “Our mission at Amgen is to serve patients by advancing medicines that address serious disease. Kyprolis is an important building block in our robust, differentiated pipeline.”

Amgen Announces Phase 3 ASPIRE Trial of Kyprolis® in Patients with Relapsed Multiple Myeloma Met Primary Endpoint

Page 2

Bradway further explained, “Coupled with our recent U.S. regulatory submissions for ivabradine and talimogene laherparepvec and our upcoming regulatory submissions for evolocumab and binatumomab, our pipeline continues to show notable progress.”

Results from the ASPIRE study will form the basis for regulatory submissions throughout the world beginning in the first half of 2015. In the U.S., the data may support the conversion of accelerated approval to full approval and expand the current indication.

“In the treatment of patients with multiple myeloma, periods of remission become shorter following each treatment regimen, underscoring the need for new options. The results of the ASPIRE study demonstrate that Kyprolis can significantly extend the time patients live without their disease progressing,” said Pablo J. Cagnoni, M.D., president, Onyx Pharmaceuticals, Inc. “The ability of novel therapies to produce deep and durable responses may, one day, transform this uniformly fatal disease to one that is chronic and manageable.”

Onyx conducted the ASPIRE study under a Special Protocol Assessment (SPA) from the U.S. Food and Drug Administration (FDA) and has received Scientific Advice from the European Medicines Agency (EMA) on the design and planned analysis of the study.

About ASPIRE

The international, randomized Phase 3 ASPIRE (CARfilzomib, Lenalidomide, and Dexamethasone versus Lenalidomide and Dexamethasone for the treatment of Patients with Relapsed Multiple Myeloma) trial evaluated Kyprolis in combination with lenalidomide and low-dose dexamethasone, versus lenalidomide and low-dose dexamethasone alone, in patients with relapsed multiple myeloma following treatment with one to three prior regimens. The primary endpoint of the trial was progression-free survival, defined as the time from treatment initiation to disease progression or death. Secondary endpoints included overall survival, overall response rate, duration of response, disease control rate, health-related quality of life, and safety. Patients were randomized to receive Kyprolis (20mg/m² on days 1 and 2 of cycle 1 only, then 27mg/m² subsequently), in addition to a standard dosing schedule of lenalidomide (25mg per day for 21 days on, 7 days off) and low-dose dexamethasone (40mg per week in 4 week cycles), versus lenalidomide and low-dose dexamethasone alone. The study randomized 792 patients at sites in North America, Europe, and Israel.

About Multiple Myeloma

Multiple myeloma is the second most common hematologic cancer and results from an abnormality of plasma cells, usually in the bone marrow. In the U.S., approximately 70,000 people are living with multiple myeloma and approximately 24,000 new cases are diagnosed annually.¹ Worldwide, nearly 230,000 people are living with multiple myeloma and approximately 114,000 new cases are diagnosed annually.² In Europe, approximately 89,000 people are living with multiple myeloma, and approximately 39,000 new cases are diagnosed annually.³

About Kyprolis® (carfilzomib) for Injection

On July 20, 2012, the FDA granted accelerated approval of Kyprolis® (carfilzomib) for Injection for the treatment of patients with multiple myeloma who have received at least two prior therapies including bortezomib and an immunomodulatory agent (IMiD), and have demonstrated disease progression on or within 60 days of completion of the last therapy. Approval was based

Amgen Announces Phase 3 ASPIRE Trial of Kyprolis® in Patients with Relapsed Multiple Myeloma Met Primary Endpoint

Page 3

on response rate. Clinical benefit, such as improvement in survival or symptoms, has not been verified.

Kyprolis is marketed in the U.S. by Onyx Pharmaceuticals, Inc., an Amgen subsidiary.

Important Safety Information Regarding Kyprolis® (carfilzomib) for Injection

Safety data have been evaluated in 526 patients with relapsed and/or refractory multiple myeloma who received single-agent Kyprolis. There were 37 deaths in the Phase 2 studies, or 7% of patients. The most common causes of death, other than disease progression, were cardiac (5 patients), end-organ failure (4 patients), and infection (4 patients). Important warnings and precautions include cardiac arrest, congestive heart failure, myocardial ischemia; pulmonary hypertension, pulmonary complications, infusion reactions, tumor lysis syndrome, thrombocytopenia, hepatic toxicity and embryo-fetal toxicity.

Death due to cardiac arrest has occurred within a day of Kyprolis administration. Patients with New York Heart Association Class III and IV heart failure, myocardial infarction in the preceding 6 months, and conduction abnormalities uncontrolled by medications were not eligible for the clinical trials. These patients may be at greater risk for cardiac complications.

Pulmonary arterial hypertension (PAH) was reported in 2% of patients treated with Kyprolis and was Grade 3 or greater in less than 1% of patients. Dyspnea was reported in 35% of patients enrolled in clinical trials. Grade 3 dyspnea occurred in 5%; no Grade 4 events, and 1 death (Grade 5) was reported.

Infusion reactions, characterized by a spectrum of systemic symptoms including fever, chills, arthralgia, myalgia, facial flushing, facial edema, vomiting, weakness, shortness of breath, hypotension, syncope, chest tightness, or angina can occur immediately following or up to 24 hours after administration of Kyprolis. Administration of dexamethasone prior to Kyprolis reduces the incidence and severity of reactions. Tumor lysis syndrome (TLS) occurred following Kyprolis administration in < 1% of patients. Patients with multiple myeloma and a high tumor burden should be considered to be at greater risk for TLS.

Thrombocytopenia following Kyprolis administration resulted in a dose reduction in 1% of patients and discontinuation of treatment with Kyprolis in < 1% of patients.

Cases of hepatic failure, including fatal cases, have been reported (< 1%). Kyprolis can cause elevations of serum transaminases and bilirubin.

There are no adequate and well-controlled studies in pregnant women using Kyprolis. Females of reproductive potential should be advised to avoid becoming pregnant while being treated with Kyprolis.

The most common serious adverse reactions were pneumonia, acute renal failure, pyrexia, and congestive heart failure. The most common adverse reactions (incidence of 30% or greater) observed in clinical trials of patients with multiple myeloma were fatigue, anemia, nausea, thrombocytopenia, dyspnea, diarrhea, and pyrexia. Serious adverse reactions were reported in 45% of patients.

Full prescribing information is available at <http://www.kyprolis.com>.

Amgen Announces Phase 3 ASPIRE Trial of Kyprolis® in Patients with Relapsed Multiple Myeloma Met Primary Endpoint

Page 4

About Amgen

Amgen is committed to unlocking the potential of biology for patients suffering from serious illnesses by discovering, developing, manufacturing and delivering innovative human therapeutics. This approach begins by using tools like advanced human genetics to unravel the complexities of disease and understand the fundamentals of human biology.

Amgen focuses on areas of high unmet medical need and leverages its biologics manufacturing expertise to strive for solutions that improve health outcomes and dramatically improve people's lives. A biotechnology pioneer since 1980, Amgen has grown to be the world's largest independent biotechnology company, has reached millions of patients around the world and is developing a pipeline of medicines with breakaway potential.

For more information, visit www.amgen.com and follow us on www.twitter.com/amgen.

About Onyx Pharmaceuticals, Inc.

Based in South San Francisco, California, Onyx Pharmaceuticals, Inc., an Amgen subsidiary, is a global biopharmaceutical company engaged in the development and commercialization of innovative therapies for improving the lives of people with cancer. The company is focused on developing novel medicines that target key molecular pathways. For more information about Onyx, visit the company's website at www.onyx.com. Onyx Pharmaceuticals is on Twitter. Sign up to follow our Twitter feed @OnyxPharm at <http://twitter.com/OnyxPharm>.

REVLIMID® is a registered trademark of Celgene Corporation.

Forward-Looking Statements

This news release contains forward-looking statements that are based on the current expectations and beliefs of Amgen Inc. and its subsidiaries (Amgen) and are subject to a number of risks, uncertainties and assumptions that could cause actual results to differ materially from those described. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including estimates of revenues, operating margins, capital expenditures, cash, other financial metrics, expected legal, arbitration, political, regulatory or clinical results or practices, customer and prescriber patterns or practices, reimbursement activities and outcomes and other such estimates and results. Forward-looking statements involve significant risks and uncertainties, including those discussed below and more fully described in the Securities and Exchange Commission (SEC) reports filed by Amgen Inc., including Amgen Inc.'s most recent annual report on Form 10-K and any subsequent periodic reports on Form 10-Q and Form 8-K. Please refer to Amgen Inc.'s most recent Forms 10-K, 10-Q and 8-K for additional information on the uncertainties and risk factors related to Amgen's business. Unless otherwise noted, Amgen is providing this information as of Aug. 4, 2014 and expressly disclaims any duty to update information contained in this news release.

No forward-looking statement can be guaranteed and actual results may differ material from those Amgen projects. Discovery or identification of new product candidates or development of new indications for existing products cannot be guaranteed and movement from concept to product is uncertain; consequently, there can be no guarantee that any particular product candidate or development of a new indication for an existing product will be successful and become a commercial product. Further, preclinical results do not guarantee safe and effective

Amgen Announces Phase 3 ASPIRE Trial of Kyprolis® in Patients with Relapsed Multiple Myeloma Met Primary Endpoint

Page 5

performance of product candidates in humans. The complexity of the human body cannot be perfectly, or sometimes, even adequately modeled by computer or cell culture systems or animal models. The length of time that it takes for Amgen and its partners to complete clinical trials and obtain regulatory approval for product marketing has in the past varied and Amgen expects similar variability in the future. Amgen develops product candidates internally and through licensing collaborations, partnerships and joint ventures. Product candidates that are derived from relationships may be subject to disputes between the parties or may prove to be not as effective or as safe as Amgen may have believed at the time of entering into such relationship. Also, Amgen or others could identify safety, side effects or manufacturing problems with Amgen's products after they are on the market. Amgen's business may be impacted by government investigations, litigation and product liability claims. If Amgen fails to meet the compliance obligations in the corporate integrity agreement between Amgen and the U.S. government, Amgen could become subject to significant sanctions. Amgen depends on third parties for a significant portion of its manufacturing capacity for the supply of certain of its current and future products and limits on supply may constrain sales of certain of its current products and product candidate development.

In addition, sales of Amgen's products (including products of Amgen's wholly-owned subsidiaries) are affected by the reimbursement policies imposed by third-party payers, including governments, private insurance plans and managed care providers and may be affected by regulatory, clinical and guideline developments and domestic and international trends toward managed care and healthcare cost containment as well as U.S. legislation affecting pharmaceutical pricing and reimbursement. Government and others' regulations and reimbursement policies may affect the development, usage and pricing of Amgen's products. In addition, Amgen competes with other companies with respect to some of its marketed products as well as for the discovery and development of new products. Amgen believes that some of its newer products, product candidates or new indications for existing products, may face competition when and as they are approved and marketed. Amgen's products may compete against products that have lower prices, established reimbursement, superior performance, are easier to administer, or that are otherwise competitive with its products. In addition, while Amgen and its partners routinely obtain patents for their products and technology, the protection of Amgen's products offered by patents and patent applications may be challenged, invalidated or circumvented by its competitors and there can be no guarantee of Amgen's or its partners' ability to obtain or maintain patent protection for Amgen's products or product candidates. Amgen cannot guarantee that it will be able to produce commercially successful products or maintain the commercial success of its existing products. Amgen's stock price may be affected by actual or perceived market opportunity, competitive position, and success or failure of its products or product candidates. Further, the discovery of significant problems with a product similar to one of Amgen's products that implicate an entire class of products could have a material adverse effect on sales of the affected products and on Amgen's business and results of operations. Amgen's efforts to integrate the operations of companies it has acquired may not be successful. Cost saving initiatives may result in Amgen incurring impairment or other related charges on its assets. Amgen may experience difficulties, delays or unexpected costs and not achieve anticipated cost savings from its recently announced restructuring plans. Amgen's business performance could affect or limit the ability of Amgen's Board of Directors to declare a dividend or their ability to pay a dividend or repurchase Amgen common stock.

Amgen Announces Phase 3 ASPIRE Trial of Kyprolis® in Patients with Relapsed Multiple Myeloma Met Primary Endpoint
Page 6

The scientific information discussed in this news release relating to new indications for Amgen's products is preliminary and investigative and is not part of the labeling approved by the U.S. Food and Drug Administration (FDA) for the products. The products are not approved for the investigational use(s) discussed in this news release, and no conclusions can or should be drawn regarding the safety or effectiveness of the products for these uses.

#

CONTACT: Amgen, Thousand Oaks
Kristen Davis, 805-447-3008 (Amgen media)
Danielle Bertrand, 650-266-2114 (Onyx media)
Arvind Sood, 805-447-1060 (investors)

¹ National Cancer Institute. SEER Stat Fact Sheets: Myeloma. Available at <http://seer.cancer.gov/statfacts/html/mulmy.html>.

² International Agency for Research on Cancer, GLOBOCAN 2012 database. Available at <http://globocan.iarc.fr/>.

³ Cancer Research UK. "Myeloma Incidence Statistics." Available at: <http://www.cancerresearchuk.org/cancer-info/cancerstats/types/myeloma/incidence/uk-multiple-myeloma-incidence-statistics#europeandworldwide>.